



Original Effective Date: 11/01/2018
Current Effective Date: 10/09/2025
Last P&T Approval/Version: 07/30/2025
Next Review Due By: 07/2026
Policy Number: C15367-A

Signifor (pasireotide diaspertate)

PRODUCTS AFFECTED

Signifor (pasireotide diaspertate), Signifor LAR (pasireotide)

COVERAGE POLICY

Coverage for services, procedures, medical devices and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Coverage Guideline must be read in its entirety to determine coverage eligibility, if any. This Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide Molina Healthcare complete medical rationale when requesting any exceptions to these guidelines.

Documentation Requirements:

Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

DIAGNOSIS:

Cushing's Disease, Acromegaly

REQUIRED MEDICAL INFORMATION:

This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. If a drug within this policy receives an updated FDA label within the last 180 days, medical necessity for the member will be reviewed using the updated FDA label information along with state and federal requirements, benefit being administered and formulary preferencing. Coverage will be determined on a case-by case basis until the criteria can be updated through Molina Healthcare, Inc. clinical governance. Additional information may be required on a case-by-case basis to allow for adequate review. When the requested drug product for coverage is dosed by weight, body surface area or other member specific measurement, this data element is required as part of the medical necessity review. The Pharmacy and Therapeutics Committee has determined that the drug benefit shall be a mandatory generic and that generic drugs will be dispensed whenever available.

A. CUSHING'S DISEASE:

1. Documented diagnosis of Cushing's Disease
- AND

Molina Healthcare, Inc. confidential and proprietary © 2025

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

Drug and Biologic Coverage Criteria

2. Documentation that pituitary surgery was not curative for the member OR member is not a candidate for pituitary surgery
AND
3. Documentation of baseline elevated urinary free cortisol
AND
4. Documentation of trial (12 weeks) and failure, serious side effects, or labeled contraindication to ketoconazole therapy

B. ACROMEGALY (SIGNIFOR LAR ONLY):

1. Documented diagnosis of acromegaly
AND
2. Documentation that member is not eligible for pituitary surgery or has had an inadequate response to pituitary surgery or radiation)
AND
3. Documentation of trial and failure, serious side effects, or contraindication to cabergoline at maximally tolerated doses for members with modest elevation of IGF-1 (defined as less than 1.4 mcg/L) and mild signs and symptoms of GH excess (2014 Guidelines) and octreotide/lanreotide

CONTINUATION OF THERAPY:

A. CUSHING'S DISEASE:

1. Adherence to therapy at least 85% of the time as verified by the prescriber or member medication fill history OR adherence less than 85% of the time due to the need for surgery or treatment of an infection, causing temporary discontinuation
AND
2. Documentation member has had a decrease in urinary free cortisol or serum cortisol from baseline levels
AND
3. Prescriber attests to or clinical reviewer has found no evidence of intolerable adverse effects or drug toxicity

B. ACROMEGALY (SIGNIFOR LAR ONLY):

1. Adherence to therapy at least 85% of the time as verified by the prescriber or member medication fill history OR adherence less than 85% of the time due to the need for surgery or treatment of an infection, causing temporary discontinuation
AND
2. (a) Documentation of growth hormone level <5 ng/mL OR IGF-1 level <1.9 U/mL for males or <2.2 U/mL for females
OR
(b) Documentation of clinical improvement (e.g., reduction in tumor size, decreased headaches, improved cardiovascular or respiratory symptoms)
AND
3. Prescriber attests to or clinical reviewer has found no evidence of intolerable adverse effects or drug toxicity

DURATION OF APPROVAL:

Initial authorization: 12 months, Continuation of Therapy: 12 months

PRESCRIBER REQUIREMENTS:

Prescribed by or in consultation with a board-certified endocrinologist or physician who specializes in the treatment of Cushing's disease [If prescribed in consultation, consultation notes must be submitted with initial request and reauthorization requests]

AGE RESTRICTIONS:

18 years of age and older

Molina Healthcare, Inc. confidential and proprietary © 2025

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

Drug and Biologic Coverage Criteria

QUANTITY:

SIGNIFOR: 0.9mg twice a day

SIGNIFOR LAR: Acromegaly: Maximum 60mg once every 4 weeks (28 days); Cushing's Disease: Maximum 40mg once every 4 weeks (28 days)

PLACE OF ADMINISTRATION:

Signifor (pasireotide diaspertate): The recommendation is that injectable medications in this policy will be for pharmacy or medical benefit coverage and the subcutaneous injectable products administered in a place of service that is a non-hospital facility-based location.

Signifor (pasireotide) LAR-- intramuscular: The recommendation is that injectable medications in this policy will be for pharmacy or medical benefit coverage and the intramuscular injectable products administered in a place of service that is a non-hospital facility-based location.

DRUG INFORMATION

ROUTE OF ADMINISTRATION:

Subcutaneous, Intramuscular (LAR)

DRUG CLASS:

Somatostatic Agents

FDA-APPROVED USES:

Signifor (pasireotide diaspertate): Indicated for the treatment of adult patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative

Signifor LAR (pasireotide): Indicated for the treatment of patients with acromegaly who have had an inadequate response to surgery and/or for whom surgery is not an option, and for patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative.

COMPENDIAL APPROVED OFF-LABELED USES:

None

APPENDIX

APPENDIX:

None

BACKGROUND AND OTHER CONSIDERATIONS

BACKGROUND:

Cushing's disease

Cushing's disease is caused by pituitary adrenocorticotrophic hormone-secreting tumors. First-line treatment is targeted at removing or destroying the tumor either by a transsphenoidal surgery or pituitary radiation therapy.

Cushing's syndrome

Cushing's syndrome is a manifestation of hypercortisolism, which can be secondary to a number of sources, chiefly, an ACTH-secreting pituitary tumor, a non-pituitary or "ectopic" ACTH-secreting tumor, or an adrenal adenoma or carcinoma that produces cortisol. Primary treatment is surgical, but when it is ineffective or cannot be performed, medical therapy is indicated.

Pasireotide is not recommended to treat other types of Cushing's Syndrome. The NCCN Guidelines

Drug and Biologic Coverage Criteria

Version 3.2018 - Neuroendocrine and Adrenal Tumors section Evaluation and Treatment of Cushing's Syndrome makes no mention for the use of pasireotide in Cushing's Syndrome caused by non- pituitary tumors. "Medical management of hypercortisolism is achieved with adrenostatic agents, including ketoconazole, mitotane, and/or mifepristone. Ketoconazole is most commonly used (at doses of 400-1200 mg/d) because of its easy availability and relatively tolerable toxicity profile. The data supporting use of other individual drugs for the management of Cushing's disease are limited. Current NCCN guidelines reference using the Endocrine Society Clinical Practice Guidelines for the Treatment of Cushing Syndrome.

Octreotide or lanreotide can also be considered for ectopic Cushing's syndrome if the tumor is somatostatin scintigraphy-positive, although it may be less effective in controlling ectopic ACTH secretion than it is in other contexts. Bilateral adrenalectomy is generally recommended when medical management of ectopic Cushing's syndrome fails."

CONTRAINDICATIONS/EXCLUSIONS/DISCONTINUATION:

All other uses of SIGNIFOR and SIGNIFOR LAR are considered experimental/investigational and therefore, will follow Molina's Off- Label policy. Contraindications to pasireotide include: avoid use in patients with severe hepatic impairment (i.e., Child-Pugh Class C).

Exclusions/Discontinuation:

Inconclusive or Non-Supportive Evidence

For neuroendocrine tumors, evidence is insufficient, conflicting, or poor and demonstrates an incomplete assessment of net benefit vs harm; additional research is recommended. (RG B) Pasireotide is a promising somatostatin analogue for the treatment of epithelial neoplasms that undergo neuroendocrine differentiation because it binds with high affinity to somatostatin receptors 1, 2, 3, and 5. Pasireotide may have a role in the treatment of octreotide-resistant carcinoid tumors. In a phase III, multicenter, randomized controlled trial, 110 adults with metastatic neuroendocrine digestive tract tumors and inadequately controlled carcinoid symptoms were assigned to administration of either long-acting octreotide or long-acting pasireotide. After 6 months, the study was halted because carcinoid symptom control appeared to be comparable between the 2 therapies; however, progression-free survival was superior for those treated with pasireotide (12 months vs 7 months). A phase II open-label multicenter study of members with advanced neuroendocrine tumors and carcinoid syndrome whose symptoms were inadequately controlled by octreotide showed that, of 44 members evaluated for efficacy, 27% had complete or partial symptom control over 15 days of treatment with pasireotide. A phase II open-label study of 29 members with metastatic neuroendocrine tumors found that pasireotide administration was associated with median progression-free survival of 11 months, although 79% of members had significant hyperglycemia.

Case reports and series have suggested possible utility of pasireotide for insulinoma. Review articles have stated that further studies are needed to better define the role, if any, of pasireotide for therapy of neuroendocrine tumors.

For postoperative pancreatic fistula prevention, evidence is insufficient, conflicting, or poor and demonstrates an incomplete assessment of net benefit vs harm; additional research is recommended. (RG B) A randomized study of 300 members undergoing pancreaticoduodenectomy or distal pancreatectomy assigned members to 7 days of pasireotide or placebo on the morning of surgery. Pasireotide administration was associated with significantly lower occurrence of pancreatic leak, fistula, or abscess as compared with placebo (9% vs 21%, respectively); significant hyperglycemia and dose-limiting nausea were higher in those receiving pasireotide.

Do not use in patients with cholelithiasis; poorly controlled diabetes mellitus (HbA1c 8% or greater at time pasireotide is initiated); or elevated QTc interval (500 milliseconds or greater).

OTHER SPECIAL CONSIDERATIONS:

Recommended treatment for hypercortisolism (Cushing's Syndrome) (Nieman) – Recommended approaches to treatment include surgical removal of the pituitary corticotropin- secreting tumor, radiation therapy with or

Drug and Biologic Coverage Criteria

without medical therapy, or medical therapy when surgery and or radiation have not been effective, or adrenalectomy (surgical or medical with mitotane). Medical therapy may include adrenal enzyme blockers (ketoconazole and/or metopirone), adrenolytic agents (mitotane), pituitary agents (cabergoline, pasireotide), or glucocorticoid receptor antagonists (mifepristone).

Recommended medical treatments and dosing

Cabergoline - Cushing syndrome (off-label use): Oral: Initial: 0.5 mg once weekly or 1 mg weekly (given as 0.5 mg twice weekly); may increase by 0.5 to 1 mg weekly at 1- or 2-month intervals until complete and sustained normalization of urinary free cortisol (UFC) levels; maximum: 7 mg weekly (given as 1 mg once daily) (ES [Neiman 2015]; Godbout 2010; Pivonello 2009)

Ketoconazole - Cushing syndrome (off-label use): Oral: Initial: 400 to 600 mg daily in 2 or 3 divided doses; may increase dose by 200 mg daily every 7 to 28 days up to a maximum of 1,200 mg daily in 2 or 3 divided doses; dosage range: 200 to 1,200 mg daily; mean effective dose in most studies: 600 to 800 mg daily in 2 divided doses (Castinetti 2014; ES [Nieman 2015]; Miller 1993)

Metopirone - Cushing syndrome (off-label use): Oral: Initial: 250 mg 4 times a day; dosage range: 500 mg to 6,000 mg daily in divided doses every 6 to 8 hours; maximum daily dose: 6,000 mg (Biller 2008; ES [Neiman 2015])

Mitotane - Cushing syndrome (off-label use): Oral: Initial: 500 mg 3 times daily (Biller 2008); may increase dose rapidly during the first 4 to 6 weeks up to a maximum of 4,000 mg to 8,000 mg per day in 3 divided doses, with the largest dose given in the evening to minimize discomfort (Baudry 2012; ES [Neiman 2015]; Schteingart 1980); after achieving control of cortisol secretion, gradually taper to the minimal dose required to maintain remission (Baudry 2012)

Mifepristone – Hyperglycemia in members with Cushing syndrome (Korlym): Oral: Initial dose: 300 mg once daily. Dose may be increased in 300 mg increments at intervals of ≥ 2 to 4 weeks based on tolerability and symptom control. Maximum dose: 1,200 mg once daily, not to exceed 20 mg/kg/day. If treatment is interrupted, reinitiate at 300 mg daily or a dose lower than the dose that caused the treatment to be stopped if interruption due to adverse reactions. Dosage adjustment of Korlym in members already being treated with strong CYP3A inhibitor therapy: Initial: 300 mg once daily; may increase dose as clinically indicated (maximum dose: 600 mg/day). Dosage adjustment of Korlym in members who require initiation of strong CYP3A inhibitor therapy: If current dose is 300 mg/day: No dosage adjustment necessary.

If current dose is 600 mg/day: Reduce dose to 300 mg once daily; if clinically indicated may increase dose to a maximum of 600 mg once daily.

If current dose is 900 or 1200 mg/day: Reduce dose to 600 mg once daily.

Note: Pasireotide is listed as a step therapy option in the Molina Clinical Policy for KORLYM- mifepristone for treating hyperglycemia secondary to Cushing's syndrome.

Recommended treatment for Acromegaly (Melmed and Katznelson) –

Surgical removal of the pituitary growth hormone-secreting adenoma is recommended as first-line unless surgery is declined, the member is a poor surgical candidate, or it is anticipated that the adenoma is not fully resectable. Surgical debulking for macroadenomas close to the chiasm and followed by medical therapy. Medical therapy options include bromocriptine, octreotide, lanreotide, pegvisomant, or abergoline. In the setting of residual disease after surgery, medical therapy is indicated. Radiation therapy is recommended if medical therapy has been ineffective or not was tolerated.

Recommended medical treatments and dosing –

Octreotide – Acromegaly: SubQ, IV: Initial: 50 mcg 3 times/day; titrate to achieve growth hormone levels < 5 ng/mL or IGF-I (somatomedin C) levels < 1.9 units/mL in males and < 2.2 units/mL in females. Usual effective dose: 100 mcg 3 times/day; range: 300 to 1,500 mcg/day. Doses above 300 mcg/day rarely result in additional benefit; if increased dose fails to provide additional benefit, the dose should be reduced. Note: Should be withdrawn yearly for a 4-week interval (8 weeks for depot injection) in members who have received irradiation. Resume if levels increase and signs/symptoms recur. IM depot injection: Members must be stabilized on subcutaneous octreotide for at least 2 weeks before switching to the long-acting depot. Upon switch: 20 mg IM intragluteally every 4 weeks for 3 months, then the dose may be modified based upon response.

Lanreotide – Acromegaly: SubQ: Initial dose: 90 mg once every 4 weeks for 3 months; after initial 3 months, continue monitoring and adjust dose as necessary based on clinical response of member, growth hormone (GH) levels, and/or insulin-like growth factor 1 (IGF-1) levels. **Pegvisomant – Acromegaly:** SubQ:

Drug and Biologic Coverage Criteria

Initial loading dose: 40 mg; maintenance dose: 10 mg once daily following initial loading dose; doses may be adjusted by 5 mg increments or decrements in 4- to 6- week intervals based on IGF-I concentrations (maximum maintenance dose: 30 mg daily) Cabergoline – Acromegaly (off-label use): The initial dose of cabergoline should be 0.5 mg once a week or 0.25 mg twice a week. The dose should be increased, if necessary, to 1 mg twice a week. Higher doses are not likely to decrease GH further. The presence of hyperprolactinemia does not consistently predict GH and IGF- 1 response.

Bromocriptine - Acromegaly: Oral: Initial: 1.25 to 2.5 mg daily increasing by 1.25 to 2.5 mg daily as necessary every 3 to 7 days; usual dose: 20 to 30 mg daily (maximum: 100 mg/day)

Endocrinology Society Guidelines –

5.1 We recommend medical therapy in a member with persistent disease following surgery. (1|QQQQ)

5.2 In a member with significant disease (i.e., with moderate-to-severe signs and symptoms of GH excess and without local mass effects), we suggest use of either a SRL or pegvisomant as the initial adjuvant medical therapy. (2|QQEE)

5.3 In a member with only modest elevations of serum IGF-1 and mild signs and symptoms of GH excess, we suggest a trial of a dopamine agonist, usually cabergoline, as the initial adjuvant medical therapy. (2|QQEE)

CODING/BILLING INFORMATION

CODING DISCLAIMER. Codes listed in this policy are for reference purposes only and may not be all-inclusive or applicable for every state or line of business. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement. Listing of a service or device code in this policy does not guarantee coverage. Coverage is determined by the benefit document. Molina adheres to Current Procedural Terminology (CPT®), a registered trademark of the American Medical Association (AMA). All CPT codes and descriptions are copyrighted by the AMA; this information is included for informational purposes only. Providers and facilities are expected to utilize industry-standard coding practices for all submissions. Molina has the right to reject/deny the claim and recover claim payment(s) if it is determined it is not billed appropriately or not a covered benefit. Molina reserves the right to revise this policy as needed.

HCPCS CODE	DESCRIPTION
J3490	Unclassified drugs (Signifor, pasireotide diaspartate)
J2502	Injection, pasireotide long acting, 1 mg

AVAILABLE DOSAGE FORMS:

Signifor SOLN 0.3MG/ML single-dose ampule

Signifor SOLN 0.6MG/ML single-dose ampule

Signifor SOLN 0.9MG/ML single-dose ampule

Signifor LAR SRER 10MG single-use kit

Signifor LAR SRER 20MG single-use kit

Signifor LAR SRER 30MG single-use kit

Signifor LAR SRER 40MG single-use kit

Signifor LAR SRER 60MG single-use kit

REFERENCES

1. Signifor LAR (pasireotide) for injectable suspension, for intramuscular use [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; July 2024.
2. Signifor (pasireotide) injection, for subcutaneous use [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; July 2024.
3. Allen PJ, Gonen M, Brennan MF, et al: Pasireotide for Postoperative Pancreatic Fistula. N Engl J Med

Molina Healthcare, Inc. confidential and proprietary © 2025

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

Drug and Biologic Coverage Criteria

2014; 370(21):2014-2022.

4. Abs R et al. Cabergoline in the treatment of acromegaly: a study in 64 members. *J Clin Endocrinol Metab.* 1998;83(2):374.
5. Baudry C, Coste J, Bou Khalil R, et al. Efficiency and tolerance of mitotane in Cushing's disease in 76 members from a single center. *Eur J Endocrinol.* 2012;167(4):473-481.
6. Biller BM, Grossman AB, Stewart PM, et al, "Treatment of Adrenocorticotropin-Dependent Cushing's Syndrome: A Consensus Statement," *J Clin Endocrinol Metab*, 2008, 93(7):2454- 62.
7. Colao A et al. A 12-Month Phase 3 Study of Pasireotide in Cushing's disease. *N Engl J Med* 2012;366:914-24.
8. Colao A, Bronstein MD, Freda P, et al. Pasireotide versus octreotide in acromegaly: a head-to-head superiority study. *J Clin Endocrinol Metab* 2014; 99:791.
9. Katznelson L et al. Acromegaly: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab.* 2014 Nov;99(11):3933-51. Epub 2014 Oct 30.
10. Giustina A, Chanson P, Kleinberg D, et al. A consensus on the medical treatment of acromegaly. *Nat Rev Endocrinol.* 2014;10(4):243-248.
11. Katznelson L, Laws ER Jr, Melmed S, et al; Endocrine Society. Acromegaly: An Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2014;99:3933-3951.
12. Oberg K, Lamberts SW. Somatostatin analogues in acromegaly and gastroenteropancreatic neuroendocrine tumours: past, present and future. *Endocr Relat Cancer.* 2016;23(12):R551- 566.
13. Manjila S, Wu OC, Khan FR, et al. Pharmacological management of acromegaly: a current perspective. *Neurosurg Focus.* 2010;29(4):E14.
14. Colao A, Bronstein MD, Freda P, et al. Pasireotide versus octreotide in acromegaly: a head-to-head superiority study. *J Clin Endocrinol Metab.* 2014;99(3):791-799.
15. Gadelha MR, Bronstein MD, Brue T, et al. Pasireotide versus continued treatment with octreotide or lanreotide in members with inadequately controlled acromegaly (PAOLA): a randomized, phase 3 trial. *Lancet Diabetes Endocrinol.* 2014;2(11):875-884.
16. Sheppard M, Bronstein MD, Freda P, et al. Pasireotide LAR maintains inhibition of GH and IGF-1 in members with acromegaly for up to 25 months: results from the blinded extension phase of a randomized, double-blind, multicenter, Phase III study. *Pituitary.* 2015;18(3):385- 394.
17. Tritos NA, Biller BM. Advances in medical therapies for Cushing's syndrome. *Discov Med.* 2012;13(69):171-179.
18. Biller BMK, Grossman AB, Stewart PM, et al. Treatment of adrenocorticotropin-dependent Cushing's syndrome: A consensus statement. *J Clin Endocrinol Metab.*2008;93:2454-2462.
19. Arnaldi G and Boscaro M. New treatment guidelines on Cushing's disease. *F1000 Med Rep.* 2009;1.
20. Mazziotti G, Gazzaruso C and Giustina A. Diabetes in Cushing syndrome: basic and clinical aspects. *Trends Endocrinol Metab.* 2011;22(12):499-506.
21. Rizk A, Honegger J, Milian M and Psaras T. Treatment options in Cushing's disease. *Clin Med Insights Oncol.* 2012(6):75-84.
22. Pavel M, O'Toole D, Costa F, et al: ENETS consensus guidelines update for the management of distant metastatic disease of intestinal, pancreatic, bronchial neuroendocrine neoplasms (NEN) and NEN of unknown primary site. *Neuroendocrinology* 2016; 103(2):172- 185.
23. Fleseriu, M., Auchus, R., Bancos, I., Ben-Shlomo, A., Bertherat, J., Biermasz, N. R., ... Biller, B. M. (2021). Consensus on diagnosis and management of Cushing's Disease: A Guideline Update. *The Lancet Diabetes & Endocrinology*, 9(12), 847–875. doi:10.1016/s2213-8587(21)00235-7

SUMMARY OF REVIEW/REVISIONS

DATE

Drug and Biologic Coverage Criteria

<p>REVISION- Notable revisions: Required Medical Information Continuation of Therapy Duration of Approval Contraindications/Exclusions/Discontinuation</p>	<p>Q3 2025</p>
<p>REVISION- Notable revisions: Required Medical Information Continuation of Therapy Quantity Background Contraindications/Exclusions/Discontinuation Available Dosage Forms References</p>	<p>Q3 2024</p>
<p>REVISION- Notable revisions: Diagnosis Required Medical Information Continuation of Therapy Duration of Approval Prescriber Requirements Age Restrictions Quantity FDA-Approved Uses Contraindications/Exclusions/Discontinuation Available Dosage Forms References</p>	<p>Q3 2023</p>
<p>REVISION- Notable revisions: Required Medical Information Continuation of Therapy Place of Administration Coding/Billing Information</p>	<p>Q3 2022</p>
<p>Q2 2022 Established tracking in new format</p>	<p>Historical changes on file</p>