



Molina Healthcare of Washington
Medicaid Private Duty Nursing
Prior Authorization Request Form
Phone Number: (800) 869-7175
Fax Number: (800) 767-7188

MEMBER INFORMATION				
Plan:	<input type="checkbox"/> Molina Medicaid (If Molina is secondary, please include a copy of the denial from primary insurance)			
Member Name:			DOB:	/ /
Member ID#:			Phone:	() -
Service Type:	<input type="checkbox"/> Elective/Routine <input type="checkbox"/> Expedited/Urgent			
REFERRAL/SERVICE TYPE REQUESTED				
Diagnosis Code & Description:				
CPT/HCPC Code & Description:				
CPT/HCPC Code & Description:				
90 DOS SPAN ONLY For continuation requests, the start date is always the day after the last authorization ends		DOS from:	/ /	
		to	/ /	
PROVIDER INFORMATION				
Requesting Provider Name:		NPI#:		TIN#:
Servicing Provider or Facility:		NPI#:		TIN#:
Contact at Requesting Provider's Office:				
Phone Number:	() -	Fax Number:	() -	
CLINICAL DOCUMENTATION TO SUPPORT NEED FOR PRIVATE DUTY NURSING (PDN)				
Signed and dated physician order for PDN [Please submit: Home Health Certification and Plan of Care, Department of Health and Human Services, HCFA Form: OBM 0938-0357]			<input type="checkbox"/> Submitted	
Current history and physical (recent hospital admissions/discharge summaries)			<input type="checkbox"/> Submitted	<input type="checkbox"/> Not Submitted
Current treatment plan and treatment records			<input type="checkbox"/> Submitted	<input type="checkbox"/> Not Submitted
Current nursing care plan - Most recent notes (two weeks)			<input type="checkbox"/> Submitted	<input type="checkbox"/> Not Submitted
Recent daily nursing notes			<input type="checkbox"/> Submitted	<input type="checkbox"/> Not Submitted
Emergency medical plan			<input type="checkbox"/> Submitted	<input type="checkbox"/> Not Submitted
90 DAY SUMMARY/including changes			<input type="checkbox"/> Submitted	<input type="checkbox"/> Not Submitted
Plan and need for more than one agency to supply care at a time?			<input type="checkbox"/> YES	<input type="checkbox"/> NO
If YES, please describe: _____				

CLINICAL PRESENTATION (check all that apply)

Frequency of assessments (to include vital signs, interventions to support patient care, health status assessment, etc.):

- Once per eight hour shift
- 2-3 times per eight hour shift
- Hourly or more often

Behavioral health, cognition, developmental monitoring:

- Non-verbal, infrequent speech, or difficult to understand
- Self-abusive behavior, risk of self-harm, and intervention required
- Sleep disturbance and patient awake more than three hours per night
- Combative, confused, or disoriented behavior that impacts self-management; patient obese
- Combative, confused, or disoriented behavior that impacts self-management

Respiratory:

- BiPAP/CPAP management
 - More than eight hours per day
 - Less than eight hours per day
- Nebulizer therapy
 - More frequent than every four hours
 - Every 4-24 hours
 - Less frequent than daily, but at least once every seven days
- Chest Physiotherapy – percussion, high-frequency chest wall oscillation vest, cough assist device, etc.
 - More than once per hour
 - Every 1-4 hours
 - Less than every four hours, but at least daily
- Oxygen management
 - Oxygen humidification, tracheal, no ventilator
 - Oxygen needed at least weekly, based on pulse oximetry
- Suctioning
 - Tracheal suctioning at least once every two hours
 - Tracheal suctioning daily, but less than every two hours
 - Nasal or oral suctioning daily
- Tracheostomy management
 - Tracheostomy management with complications (skin breakdown, replacement needed)
 - Tracheostomy management, no complications
- Ventilator management
 - Continuous ventilator use
 - Ventilator use for 12 or more hours per day
 - Ventilator use for 7-12 hours per day
 - Ventilator use for less than seven hours per day
 - Interventions in place for active weaning
 - Ventilator weaning achieved; requires ongoing post-weaning monitoring and management
 - Ventilator on standby, respiratory assistance, or used at night for less than one hour

Skilled Nursing Needs:

Blood draw

_____ Central line

_____ Peripheral line

_____ More than twice per week

_____ Less than twice per week

Infusion therapy

Blood or blood product

Chemotherapy infusion

Central line access and management

Pain medication infusion

Intravenous Infusion (IV antibiotics, etc.), including infusion administration and monitoring for infusion reactions

Infusions more than every four hours

Infusions less than every four hours

Non-infusion medication

Insulin administration

Non-insulin medication injectable administration

Medication administration at least every two hours, requiring clinical monitoring

Activity of Daily Living (ADL)/Therapy support

_____ Bedbound

_____ Wheelchair user

_____ Ambulatory

Total/partial lift, weight 55-125 pounds

Total/partial lift, weight greater than 125 pounds

ADL support needed more than four hours per day to maximize patient's independence

Body cast management

Cast or brace management

Splinting management, including removal and replacement, at least every eight hours

Communication deficit; nurse to support therapy plan

Range of motion exercises at least every eight hours

Physical therapy program at least three hours per day; occupational therapy program at least four hours per day

Nutrition management

Enteral nutrition with complications, requires administration of feeding, residual check, adjustment or placement of tube, and assessment or management of complications

Enteral nutrition without complications

Gastrostomy tube care, uncomplicated

Nasogastric tube care, uncomplicated

Partial parenteral nutrition with central line care

Total parenteral nutrition with central line care

Skin and wound care management

Burn care

Ostomy care, at least once per day

Postsurgical care, within 45 days of surgery

Stage one or two wound management, at least once per day

Stage three or four wound management, at least once per day

Stage three or four wound management at least once per day, and multiple wound sites

Prescribed topical medication application at least every four hours

Wound vacuum management

- Seizure control that requires nursing intervention/management
 - Seizures lasting less than three minutes, at least four times per week
 - Seizures lasting 3-5 minutes, at least four times per week
 - Seizures lasting 3-5 minutes, one to four times per day
 - Seizures lasting 3-5 minutes, more than five times per day
 - Seizures lasting more than five minutes, or clustered seizures, or seizure activity without regaining consciousness, at least four times per week
 - Seizures lasting more than five minutes, or clustered seizures, or seizure activity without regaining consciousness, one time or more per day, requiring rectal medication
 - Seizures lasting more than five minutes, or clustered seizures, or seizure activity without regaining consciousness, one time or more per day, requiring IM or IV medication

ADDITIONAL INFORMATION

List: